

**MICON**


**94**

**INTERNATIONAL**

&

**35TH ANNUAL CONFERENCE OF  
ASSOCIATION OF MICROBIOLOGISTS OF INDIA**

**9 - 12 NOVEMBER 1994**



**INVITED PAPERS  
POSTER ABSTRACTS  
AND SOUVENIR**

**DEFENCE FOOD RESEARCH LABORATORY  
SIDDARTHA NAGAR, MYSORE - 570 011 INDIA**



---

EMB-88

**BIOMINERALIZATION OF  $\alpha$ - $\beta$ - $\gamma$  AND  $\delta$ -  
ISOMERS OF HEXACHLOROHEXANE BY  
MICROBIAL CONSORTIA**

N. SREEDHAR REDDY AND A.A.M. KUNHI  
Department of Microbiology and Bioengineering, Central  
Food Technological Research Institute  
Mysore - 570 013.

Detection of high levels of HCH and other organo-chlorine pesticide residues in food and human adipose tissues in India and other developing countries have been a matter of great concern. It is highly necessary to develop treatment techniques to eliminate these recalcitrant and toxic chemicals from the environment and industrial effluents. Microbial consortia capable of degrading  $\alpha$ - $\beta$ - $\gamma$  and  $\delta$ -isomers of HCH have been developed in the laboratory by long term enrichment of soil and sewage samples using different isomers of HCH (separately) as the sole source of carbon and energy. The consortia designated as AHR, BHR, GHR and DHR degraded  $\alpha$ -HCH (upto 50 ppm),  $\beta$ -HCH (10 ppm),  $\gamma$ -HCH (25 ppm) and  $\delta$ -HCH (10 ppm), respectively both under shaken and stationery conditions. In all the cases stoichiometric amounts of chloride was released indicating complete mineralization of the compounds. Addition of auxiliary carbon source such as glucose, cellulose and saw dust improved the rate of degradation of  $\alpha$ -HCH by AHR. The optimal temperature and pH were 30°C and 7.5 respectively. About 10 different bacterial strains were observed in the consortia.

---

EMB-70

**ESTIMATION OF BASIC KINETIC  
PARAMETERS IN THE MICROBIAL  
DEGRADATION OF ALPHA-  
HEXACHLOROCYCLOHEXANE**

N. SREEDHAR REDDY, M. K. GOWTHAMAN AND  
A.A.M. KUNHI  
Department of Microbiological and Bioengineering,  
Central Food Technological Research Institute  
Mysore-570 013.

Microbial degradation of  $\alpha$ -Hexachloro-cyclohexane (HCH) is of great importance in pollution control. This study deals with the determination of kinetic parameters during  $\alpha$ -HCH degradation by a microbial consortia on which no literature is available. By the application of the simple growth, Monod's and logistic models  $\mu_{max}$ ,  $K_s$ ,  $K_d$  and degradability have been determined as a preliminary approach to more complex modelling required in the future.  $\mu_{max}$  values ranged from 0.469 to 0.214 day<sup>-1</sup> for  $\alpha$ -HCH concentration range of 5.0 to 50mg/L with the corresponding  $K_s$  ranging from 1.96 to 15.88 mg/L. The substrate degradation rate  $K_d$  ranged from 0.197 to 0.326 day<sup>-1</sup> with  $\alpha$ -HCH alone, while, in the presence of glucose, cellulose and sawdust (co-substrates) it increased to 0.425, 0.413 and 0.383 day<sup>-1</sup> respectively.